

REFERENCE LIST:

2002 (14)

- Ali S.N.H., Ramanna M.S., Jacobsen E. and Visser R.G.F. 2002. Genome differentiation between *Lycopersicon esculentum* and *L-pennellii* as revealed by genomic in situ hybridization. *Euphytica* 127: 227-234.
- Appeldoorn N.J.G., Sergeeva L., Vreugdenhil D., Van der Plas L.H.W. and Visser R.G.F. 2002. In situ analysis of enzymes involved in sucrose to hexose-phosphate conversion during stolon-to-tuber transition of potato. *Physiologia Plantarum* 115: 303-310.
- Bachem C., Fernandez del Carmen A. and Visser R.G.F. 2002 cDNA-AFLP based amplified-fragment length polymorphism fingerprinting. In: Lorkowski S. and Cullen P. (eds), *Analysing Gene Expression: Wiley-VCH*. pp. 285-295.
- Bonnema G., Berg P. and Lindhout P. 2002. AFLPs mark different genomic regions compared with RFLPs: a case study in tomato. *Genome* 45: 217-221.
- Bovy A., De Vos R., Kemper M., Schijlen E., Pertejo M.A., Muir S., Collins G., Robinson S., Verhoeyen M., Hughes S., Santos-Buelga C. and Van Tunen A. 2002. High-flavonol tomatoes resulting from the heterologous expression of the maize transcription factor genes LC and C1. *Plant Cell* 14: 2509-2526.
- Brugmans B., Del Carmen A.F., Bachem C.W.B., Van Os H., Van Eck H.J. and Visser R.G.F. 2002. A novel method for the construction of genome wide transcriptome maps. *Plant Journal* 31: 211-222.
- De Wit P.J.G.M., Brandwagt B.F., Van den Burg H.A., Cai X., Van der Hoorn R.A.L., De Jong C.F., Van't Klooster J., De Kock M.J.D., Kruijt M., Lindhout W.H., Luderer R., Takken F.L.W., Westerink N., Vervoort J.J.M. and Joosten M. 2002. The molecular basis of co-evolution between *Cladosporium fulvum* and tomato. *Antonie Van Leeuwenhoek International Journal Of General And Molecular Microbiology* 81: 409-412.
- Eastmond P.J., van Dijken A.J.H., Spielman M., Kerr A., Tissier A.F., Dickinson H.G., Jones J.D.G., Smeekens S.C. and Graham I.A. 2002. Trehalose-6-phosphate synthase 1, which catalyses the first step in trehalose synthesis, is essential for *Arabidopsis* embryo maturation. *Plant Journal* 29: 225-235.
- Edwards A., Vincken J.P., Suurs L., Visser R.G.F., Zeeman S., Smith A. and Martin C. 2002. Discrete forms of amylose are synthesized by isoforms of GBSSI in pea. *Plant Cell* 14: 1767-1785.
- Heeres P., Schippers-Rozenboom M., Jacobsen E. and Visser R.G.F. 2002. Transformation of a large number of potato varieties: genotype-dependent variation in efficiency and somaclonal variability. *Euphytica* 124: 13-22.
- Horvath B.M., Bachem C.W.B., Trindade L.M., Oortwijn M.E.P. and Visser R.G.F. 2002. Expression analysis of a family of nsLTP genes tissue specifically expressed throughout the plant and during potato tuber life cycle. *Plant Physiology* 129: 1494-1506.
- Lindhout P. 2002. The perspectives of polygenic resistance in breeding for durable disease resistance. *Euphytica* 124: 217-226.

- Oomen R., Doeswijk-Voragen C.H.L., Bush M.S., Vincken J.P., Borkhardt B., van den Broek L.A.M., Corsar J., Ulvskov P., Voragen A.G.J., McCann M.C. and Visser R.G.F. 2002. In muro fragmentation of the rhamnogalacturonan I backbone in potato (*Solanum tuberosum* L.) results in a reduction and altered location of the galactan and arabinan side-chains and abnormal periderm development. *Plant Journal* 30: 403-413.
- Yuan Y.N., Haanstra J., Lindhout P. and Bonnema G. 2002. The *Cladosporium fulvum* resistance gene Cf-ECP3 is part of the Orion cluster on the short arm of tomato Chromosome 1. *Molecular Breeding* 10: 45-50.